

Review article

# Nutrition as an integral part of behavioural management of zoo animals

Anouk Fens<sup>1,2</sup> and Marcus Clauss<sup>3</sup>

<sup>1</sup>Apenheul Primate Park, Postbus 97, 7300 AB Apeldoorn, The Netherlands

<sup>2</sup>Amersfoort Zoo, Barchman Wuytierslaan 224, 3819 AC Amersfoort, The Netherlands

<sup>3</sup>Clinic for Zoo Animals, Exotic Pets and Wildlife, Vetsuisse Faculty, University of Zurich, Winterthurerstr. 260, 8057 Zurich, Switzerland

Correspondence: Marcus Clauss, email; mclauss@vetclinics.uzh.ch

**Keywords:** activity budget, diet, captivity, dietary management, feeding behaviour, mammal, nutrition

**Article history:**

Received: 17 Aug 2023

Accepted: 17 Apr 2024

Published online: 31 Oct 2024

**Abstract**

Provision of optimal diets for wild animals in captivity can have beneficial effects on health, reproductive performance and welfare. Zoo animal nutrition has made significant advances over the past decades; however adequate diet provision in zoological institutions remains challenging. This paper proposes a schematic history of developments in zoo animal nutrition in four steps, from finding diet items an animal will readily accept to secure its immediate survival (step 1), to supplementing these diet items with essential nutrients like minerals and vitamins (step 2), replacing convenient diet items from step 1 with items that are nutritionally and structurally more adequate for the species (the culturally challenging step 3) and providing the step 3 diet in species-appropriate ways to achieve behavioural management goals (step 4). Within this model, general rules of feeding behaviour are considered, emphasising the importance of promoting natural behaviours through adequate feeding management rather than simply preventing nutritional imbalances. By providing several short case studies, knowledge of feeding behaviour and activity budgets of several mammalian species is used as a guideline for adequate feeding management. Since the developments in feeding regimes described often cannot be made in synchrony across species or across zoos, it remains challenging for zoos to progress towards the final step. Therefore, this paper aims to inspire zoos by providing suggestions for moving towards integral feeding regimes for animals in their care and to assess where they stand in the four-step process with respect to certain animal groups. In conclusion, active application of knowledge of both dietary requirements and natural feeding ecology are essential for improving dietary feeding management in zoos and hence animal welfare.

## Introduction

The provision of an appropriate diet is a crucial element of animal husbandry, including of zoo animal populations. Nutrition has distinct effects on animal health, reproductive performance and welfare. When considering zoo animal nutrition, the focus of attention may vary among these aspects, or logistical aspects of availability, hygiene or costs. Although there have been considerable developments in the science of zoo animal nutrition and dietary management, providing a nutritionally balanced diet to a broad array of taxonomic groups remains challenging. This review stresses the role nutrition plays in the behavioural management of wild animals in captivity. First, a summative overview of historical developments in zoo

animal nutrition is given. Then some general rules on feeding behaviour are outlined. Several short examples illustrate how nutrition influences behaviour in human care. This paper does not aim to provide a comprehensive literature review but gives citations as examples. For brevity, the focus and examples are drawn from this taxon. Nevertheless, similar principles will apply to other taxa.

### *A brief history of zoo animal nutrition*

The general history of zoo animal nutrition and the history of developing a feeding regime for a particular species can be summarised in a four-step model. This model may not be universal and comprehensive. While the sequence appears relatively constant, it may be that while a majority of zoos are

already in step 3 for a group of species, they are still in step 2 for another set of species; and some other zoos might still be in steps 2 and 1, respectively. Developments are not made in synchrony across species, across zoos or even across species and sections within an individual zoo. Evaluating to what extent anecdotal or scientific communication affects progress from one step to another within or between zoos is beyond this review. Zoos have been sharing diet information for a very long time, for example in the 1966 issue of the International Zoo Yearbook (Hediger 1966; Ratcliffe 1966; Wackernagel 1966) or the 1977 CRC handbook series (Rechcigl 1977a, b) and in numerous individual publications and husbandry guidelines. It is expected that a recently founded zoo or a zoo that begins keeping a new species nowadays would begin the feeding management process at step 3 or 4 rather than step 2. However, the authors have encountered dietary recommendations and diets actually fed that do not correspond to current nutritional knowledge.

#### Step 1: Immediate survival

In the initial stages of zoo husbandry of a species, it is crucial to find a food that the animals will actually ingest, hence allowing their survival in the short term. This step may seem intuitive and self-explanatory. Carnivores would be offered meat. Herbivores will most likely not survive on meat diets, even if they may voluntarily ingest more animal matter than would be expected (Clauss et al. 2016). Because of the flexible nature of most animals, a complete rejection of what is chosen by keepers might be rare. An intriguing example is an experiment with nutria *Myocastor coypus*, a herbivorous rodent, that was described as having a “herbivore creed” that led to a “martyr death by self-inflicted starvation” when an individual was exposed to a non-herbivorous diet (Bickel and Geréz 1936). An important, questionable aspect of this approach is that ‘nutritional wisdom’ is ascribed to animals and hence a food item that is accepted is often considered ‘right’, even though this concept is usually rejected (Kawata 2008; Ullrey 1989). The evolutionary adaptations of animals will only lead to reasonable choices if their environment is identical to that in which the adaptations evolved—for example, in a world without cultivated fruit, grains or (in the case of humans) sweets. Additionally, in order for ‘nutritional wisdom’ to develop, consequences of a diet must be immediate so they can be related to the diet. While such a feedback mechanism may cause aversion to something bitter, it will not link the negative long-term consequences of obesity to a sweet or fatty taste.

The choice of food items at step 1 is typically from the range of foods available for consumption by humans and agricultural production animals. In composition and concept, nutritional management at this stage resembles either that for farm animals or a human diet. Because of this, the general intuitive acceptance of such diets by humans—curators, keepers, visitors—is high.

Primate diets in particular have contained typical human food items, such as when a gorilla’s *Gorilla gorilla* preferred item was sausages (Bickel and Geréz 1936). Experimental work by Remis (2002) showed that when given a choice, zoo gorillas do not choose diet items that have a nutritional composition similar to the food they ingest in the wild, but will choose items of lower fibre and higher sugar content. The parallel to human behaviour is evident. Human dietary habits persisted in zoo animal feeding for a long time, for example including dairy products in the diet of gorillas (Lukas et al. 1999) and the still widespread use of cultivated fruit for many animals that would not find food of this nutritional composition in the wild (Schmidt et al. 2005; Schwitzer et al. 2009). However, most animals will accept these items, which means animals can be kept alive for certain periods of time and may even reproduce. Rare examples of surprising reluctance include when certain herbivores cannot be sustained on grass hay,

like giraffes *Giraffa camelopardalis* ssp. and some other browsing ruminants and tapirs *Tapirus* spp. (Clauss et al. 2003; Foose 1982). But these animals accept a variety of other feeds intended for production animals or humans and can be kept alive on these foods for certain time periods. Cases of extreme specialisation that preclude basically all but a single group of diet items, such as koala *Phascolarctos cinereus* surviving only when provided with eucalyptus leaves (Hume 2005), are very rare.

Using carnivores as an example, feeding chunks of muscle meat would be the equivalent of step 1. The term ‘carnivore’ translates to ‘meat-eater’ and muscle meat resembles the meat humans acquire for their own cooking and thus is intuitively considered food. The underlying question of step 1 is: What diet will this animal accept? The welfare aspect of step 1 is immediate survival.

#### Step 2: Meeting nutritional requirements

When animals are kept on items from human or production animal diets, nutritional deficiencies may arise. A classic example is the metabolic bone disease carnivores develop when fed meat—“the original animal model of rickets” (Chesney and Hedberg 2010). Meat without bone contains too little calcium for growth, reproduction and maintenance requirements (Allen et al. 1996). The same problems occur in animals such as primates when predominantly maintained on cultivated fruit (Fiennes 1974) as fruits also contain too little calcium. Similar to the metabolic functions of calcium, insufficient vitamin D<sub>3</sub> production in primates may result in corresponding clinical signs such as rickets, growth impairment and renal failure (Crissey et al. 1999; Yamaguchi et al. 1986). Limited or non-existent ultraviolet B exposure for vitamin D<sub>3</sub> synthesis can lead to vitamin D deficiencies, in particular reported in primates with dark skin pigmentation (Moititi et al. 2022; Ziegler et al. 2018). In contrast, New World monkeys like marmosets require higher levels of dietary vitamin D (Takahashi et al. 1985). A similar logic may apply to other minerals, vitamins and protein or amino acids depending on species and feeds used. Many of these problems were addressed early on in the history of zoo animal nutrition. Herbert Ratcliffe from Philadelphia Zoo was a pioneer in this respect, introducing diets that contained nutrients in calculated concentrations to meet the requirements of specific animal groups. The effect of this change was drastic—animal survival improved with an approximate reduction in overall annual mortality of mammals and birds from 20% to 10% (Ratcliffe 1966). For example, an analysis of museum skeletons demonstrated a historical decrease in the occurrence of metabolic bone disease in zoo baboons *Papio* spp. over time (Sadhir et al. 2022).

This higher survival is also marked by an increase in the occurrence of a disease typical for old age—cancer (Lombard and Witte 1959). Better diets lead to more cancer and most likely other age-related diseases because they allow animals to reach an age where these diseases can develop (Clauss and Müller 2024). And of course, these changes also increase the likelihood of self-sustaining populations.

Thus Step 2 is characterised by a focus on mineral, vitamin and other nutrient supplementation, either in the form of supplements that are applied to deficient food items, as when meat, insects or fruit are dusted with a calcium-containing powder, or in the form of compound feeds (such as pellets, extrudates, sausages) designed as complete feeds or as part of an animal’s ration. As long as this step consists of adding supplements to the diet items chosen in step 1, it is typically well accepted by those involved because it combines the acceptance of human and agriculture-derived feeding concepts with the expertise of adding ‘a special powder’. By contrast, if this step consists of adding completely mineralised and vitaminised compound feeds to a diet, it may meet more emotional resistance because such diet items are unduly perceived as ‘artificial’ (Hediger 1966).

It is important that nutrient deficiencies are not forgotten. Classic ration calculation is challenging and best left to specialists. It aims to prevent nutrient deficiencies (or toxicities). In terms of animal biology, this step is related to the biochemistry of feeds and animals—something not intuitive for most people working in zoos.

For carnivores, feeding chunks of muscle meat dusted with a supplemental powder or feeding a mix composed of processed meat enriched with minerals and vitamins would be the equivalent of step 2. The underlying question of step 2 is, given the assumption that avoiding deficiencies will allow both reproduction and the raising of healthy offspring: What diet will allow this animal to reproduce? The welfare aspect of step 2 is freedom from deficiency-triggered disease.

#### *Step 3: Meeting physiological needs*

Success in step 2 allows focus on more subtle nutritional issues that go beyond basic survival and the prevention of dramatic deficiencies. These aspects refer to diseases or health problems that are less drastic and potentially develop over longer time periods than deficiencies. These problems are, to the authors' knowledge, classically not defined as deficiencies but often referred to as a 'lack' of a certain factor. There do not seem to be any clear-cut definitions to distinguish these. One typical example is the need for structural fibre to foster peristalsis and proper gut function in herbivores, ranging from the sheer issue of a "lack of bulk" (Meredith and Prebble 2017) to problems in colonic separation in coprophagic herbivores (Guerra Aldrigui et al. 2018) or lack of sufficient long-stemmed material for rumination in ruminants and camelids (Mertens 1997). A lack of structural components in processed meat products may lead to calculus formation on carnivore teeth (Bond and Lindburg 1990), which sometimes is countered by sporadically offering bones in addition to processed meat diets. There are potentially beneficial effects of indigestible components for various species, from the soil inadvertently ingested by insectivores (Gull et al. 2015) to the less digestible components of whole prey (Depauw et al. 2013). Finally, there are a suite of civilisation diseases with obesity at the forefront (Morfeld and Brown 2016; Schwitzer and Kaumanns 2001) and including other consequences of consuming energy-dense and often sugar-rich diets, such as malfermentation, acidosis, diabetes, hypertension and hoof problems (e.g. Kuhar et al. 2013).

Step 3 includes a move away from the diet items accepted in step 1 and towards diets that resemble, in chemical composition as well as physical structure, animals' natural diets. Often, this means a departure from long-standing food purchase traditions, requiring more engagement when the step is initiated until the new channels and procedures are well-established. Also, because diet items used in step 1 are often widely accepted, shifting away from them might be a source of social conflict among the personnel involved. As 'more natural' often means 'less palatable' compared to cultivated diet items, the animals might not react positively either.

Step 3 expands the biochemical view of nutrition towards its interactions with large anatomical structures. It is here that considerations about dentition, gastrointestinal anatomy and physiology come into play—factors that are well-known by biologists and those interested in the respective animals, including curators, keepers and informed visitors. Cultural traditions, so to speak, clash with biological logic.

It is at this step that it is recognised that cultivated fruit do not resemble the fruit that frugivores consume in natural habitats (Schmidt et al. 2005; Schwitzer et al. 2009) and that health issues related to sugar, such as obesity and dental caries, can be avoided by removing cultivated fruit from the diet (Plowman

2013). Therefore, whether a zoo is at step 2 or 3 might be assessed from which animals still receive cultivated fruit in their zoo rations. Mimicking natural diets might seem straightforward in some cases, such as providing browse for browsers and whole prey for carnivores, but is less intuitive when it comes to replacing cultivated fruit with green leafy vegetables in the diet of frugivorous primates.

For carnivores, feeding whole prey would be the equivalent of step 3. The underlying question of step 3 is: What diet will guarantee a disease-free, long life? The welfare aspect of step 3 is freedom from long-term compromised physiological health.

#### *Step 4: Meeting psychological needs*

In natural habitats, animals perform species-specific behaviours to acquire and process food which have become part of their evolutionary psychological and physiological repertoire. If food is presented in human care in ways that do not foster these behaviours to the same degree—both quantitatively (in terms of the time spent foraging and feeding) and qualitatively (in terms of the specific behaviours performed)—then behaviour at the zoo will deviate distinctively from that in the natural habitat. While time 'freed' in this way may in certain cases allow the animal to express potentially positive behaviours it cannot display in the wild, such as the novelty-seeking and innovative behaviour of orang-utans in human care (van Schaik et al. 2016), the lack of opportunity to perform a behaviour the animal has evolved for mostly leads to either excessive passivity or undesired behaviours such as stereotypies and hence compromised welfare (Bashaw et al. 2007; Britt 1998; Rees 2009). A reduction in, or absence of, undesired behaviours is an intuitive aim, the performance of species-appropriate behaviour can be interpreted as a positive affective state (Gray 2017; Mellor 2016) that is even linked to neurohormonal feedback circuits (Mellor 2015a, b) or as a means of facilitating a meaningful life for the animal, based on the fact that it is provided with opportunities to perform its natural behaviour in ways that have relevance for its life (Clauss and Schiffmann 2022). In terms of feeding, this relevance translates into a system where the animal acquires food dependent on whether it pays attention, focuses or performs work. This means that a decision by the animal to exhibit a certain behaviour has relevance and provides 'meaning', as opposed to a situation where no matter what the animal does, it will always get the food (and hence its behaviour has no meaningful effect on the outcome).

Thus, step 4 puts the focus on a holistic view of both physiological and psychological health in terms of the activity induced by the feeding regime. In other words, modern zoo feeding management addresses typical issues of energy and nutrient requirements, not only structural needs covered by diets, but also behavioural aspects. Ways of addressing this latter aspect are traditionally called 'enrichment'. Although this term is useful, it betrays a past of a less-than-optimal focus on animal welfare—because many things that make life in zoos closer to nature were called 'enrichment', as if one was doing something extra. In reality, not providing most sources of enrichment equals pauperisation, with corresponding negative impact on animal behaviour.

Many examples of discrepancies between natural habitats and zoos are easily understood. At the zoo, a predator is generally not cognitively challenged in terms of securing prey—no matter how it behaves, it will ultimately be presented with food. An elephant that receives its daily ration as a pile of hay (and pellets and vegetables) does not have to move at all once it has reached that pile, in contrast to free-ranging animals that need to move constantly while foraging. A giraffe that can dip its snout into a trough full of pellets and thus grab as large a mouthful as its lips allow will need fewer individual feeding movements than a free-ranging giraffe that must pluck individual leaves from between the

acacia's thorns.

In the authors' view, step 4 represents the most recent development in zoo animal nutrition. Whereas the problems encountered during step 3 might often be emotional, those encountered during step 4 are often logistical (in terms of the additional resources required to move from an unengaged to an engaged feeding method). Progress may also be hampered by a lack of behavioural knowledge and a lack of available solutions for practical implementation of potential methods. However, in terms of compliance, the changes necessary for step 4 are expected to meet less emotional resistance than step 3 (unless traditional husbandry methods have become ingrained) because displaying appropriate behaviours should make animals more fascinating to observe and possibly more content.

For carnivores, feeding whole prey in a way that requires cognitive as well as physical activity by the predator would be the equivalent of step 4. The underlying question of step 4 is, adopting the concept of a meaningful life from Clauss and Schiffmann (2022) outlined above: What diet and feeding method will guarantee a disease-free, long and meaningful life? The welfare aspect of step 4 is long-term physiological health with the positive or neutral affective state that comes from needing to perform adequate behaviours to have a meaningful life.

### Overview of mammalian feeding behaviour

To have aims for behavioural management, knowledge about the feeding behaviour and activity budget of the animal in question is required. This should be acquired on a species-by-species basis whenever possible. However, some rough guidelines can be outlined using a graph from Hiimae (2000) (Figure 1).

Small animals can live on comparatively small and spatially distributed items like invertebrates or nectar and find sufficient amounts of these to fulfil their daily energy requirements. Such animals typically search for their food for a large part of the day, which cannot be replicated by offering food once or twice in a bowl. As herbivores (right side of Figure 1) become larger, they have to shift their focus from items that are comparatively rare in the environment (such as seeds) to items that can be encountered everywhere (like grass or tree leaves) so that they can ingest sufficient amounts to meet their daily energy requirements. Along this axis, animals spend a large amount of their active time feeding, which can be mimicked by either feeding them repeatedly during the day—as is common practice in great apes (e.g. Bloomsmith and Lambeth 1995)—or by ensuring forage is available at all times—as is common practice in large hoofstock. As faunivores (left side of Figure 1) become larger, they may still go for comparatively small prey, such as felids hunting rodents or pinnipeds hunting fish. To meet daily energy requirements, a large number of successful individual hunts are necessary, which may be facilitated in pinnipeds by hunting prey that occurs in schools. For terrestrial predators such as a small felid living on rodents, this implies a comparatively large number of successful, individual hunts per day (De Cuyper et al. 2019), which will not be mimicked by one or two daily deliveries of food. For large terrestrial predators such as the tiger, small prey items like rodents are not accessible in sufficient numbers per day; therefore, these animals have to hunt larger prey (Carbone et al. 1999). This can often lead to a situation where these animals do not need to have a successful hunt every day but can gorge themselves to be “full and lazy” (De Cuyper et al. 2019; Jeschke 2007), a situation that is not mimicked by allotting small daily rations or by interspersing fast days in a series of days with small rations (Kleinlugtenbelt et al. 2023). Finally, large omnivores such as bears may use the opportunity offered by special ecological circumstances where they can harvest surprisingly small food items (compared to their own body size) due to the extremely high, lumped seasonal

availability of these items such as berries or salmon. Mimicking such feeding is also not feasible with one or two lumped feedings per day.

In conclusion, the feeding patterns of most mammals do not correspond to a one- or two-meal pattern that would best suit daily feeding schedules by zoo personnel. Therefore, methods to dispense food at naturalistic frequencies, ideally without increasing the workload, are required. To the authors' knowledge, corresponding feeding methods are in use for certain species in many zoos.

### Case study examples


A detailed catalogue of activity budgets and foraging patterns would be welcome for the development of guidelines for zoo animal feeding, but does not yet exist. Therefore, decisions must typically be made on a species-by-species basis or possibly on the basis of larger animal groups with similar behavioural adaptations. Several measures to achieve more naturalistic feeding behaviour are self-evident, such as distributing food across many different locations, providing food at a higher frequency than once or twice a day, making access to food more challenging both in terms of physical and cognitive effort required to attain it and choosing—from among a variety appropriate for a species—feeds with a lower energy density so that more has to be ingested for the same energy gain. Nevertheless, a variety of examples are briefly mentioned below to inspire curators and keepers to further explore the biology and solutions for mimicking natural conditions to facilitate naturalistic feeding behaviour.

#### Rabbits *Oryctolagus cuniculus*


Under natural conditions, rabbits feed for approximately 4–7 hours per day (Mykytowycz and Rowley 1958). By contrast, rabbits fed a complete laboratory rabbit diet spend approximately two hours per day feeding (Olivas et al. 2013). A study with pet rabbits showed that on a hay-only diet, rabbits fed for approximately 10 hours per day, whereas on a muesli mix diet with a higher energy density and without access to hay, the animals fed for only 2.5 hours per day (Prebble et al. 2015a). The rabbits on the muesli mix diet gained more weight and had a more obese body condition whereas all animals grew adequately, irrespective of diet (Prebble et al. 2015b). This example underlines the relevance of diet for the daily activity budget of animals. In the case of rabbits, feeding a roughage-only diet as opposed to a muesli mix or a complete laboratory diet represents a daily difference of up to 7.5 hours during which the animals are either occupied eating or have ‘free time’. Over the course of a year, this corresponds to more than 100 days of ‘nothing to do’—highlighting the relevance of nutrition for behavioural management of animals. In a professional husbandry system, there would have to be a concept of how the animals are supposed to spend their time if fed on the high-energy diet. These implications reported in rabbits can likely also contribute to appropriate dietary management in herbivorous zoo rodents such as prairie dogs *Cynomys* spp. or hyraxes *Procavia*, *Heterohyrax* and *Dendrohyrax* spp.

#### Giraffe *Giraffa camelopardalis*

Veasey et al. (1996) compared the activity budget of zoo giraffes with that of free-ranging animals. Their data are presented in Figure 2, with additional behavioural data results from other studies. It is evident that during the day (Figure 2A) zoo giraffes spend less time feeding than free-ranging giraffes and show different degrees of stereotyping. Only in a single zoo report on a feeding regime with a very high proportion of browse (Schüßler et al. 2015) were feeding times approximated that were similar to the wild and stereotypies minimised. During the night, zoo giraffe typically ruminate less than their free-ranging counterparts (Figure



**Figure 1.** Schematic representation of mammalian body size and trophic group (omnivores in the middle, faunivores on the left and herbivores on the right). The graph indicates changes of diet items with changes in body size across species, from small at the bottom centre to large towards the upper part of the graph. Modified from Hiimae (2000).



**Figure 2.** Activity budgets for giraffe *Giraffa camelopardalis* kept in zoos and in a natural habitat for (A) the daytime and (B) the nighttime (Bashaw 2011; del Castillo et al. 2005; du Toit and Yetman 2005; Orban et al. 2016; Pellew 1984; Schüßler et al. 2015; Veasey et al. 1996)

2B). These data give evidence for the difference in activity budgets of free-ranging and captive individuals and provide benchmarks for assessing interventions. As a word of caution, rather than understanding data from the natural habitat as a fixed biological characteristic, it is rather its magnitude that should serve as a guideline.

#### Equids

Free-ranging plains zebras *Equus burchelli* spend 11–15 hours grazing and 4.5–7.5 hours standing (Neuhaus and Ruckstuhl 2002). In domestic horses kept exclusively indoors, standing time (without anything to do) increased from about 8 hours when animals were fed only hay to about 13 hours when fed only pellets, with an additional increase in searching behaviour to more than 2 hours on the all-pellet diet (Elia et al. 2010). Again, this emphasises the relevance of the dietary regime on behavioural management. In a simple experiment with domestic horses, Ellis et al. (2015) increased the feeding time and reduced the standing time significantly when offering hay in a triple layer of hay nets as compared to using a single hay net. Evidently, reducing the mesh or opening size in nets, racks or similar structures will require more time of the animal to harvest a similar amount of food. Such methods should be applicable to basically all herbivores. In zoo giraffe, this principle has recently been shown to increase feeding times and concomitantly reduce stereotypies (Depauw et al. 2023; Walldén 2023).

#### Sloth *Choloepus hoffmanni*

The diet fed to two-toed sloths in zoos differs distinctively from that in the wild; in particular, many items fed in zoos are higher in energy density (Hayssen 2011). At the same time, the reported daily time spent sleeping is 20 hours at zoos versus only 11 hours in the natural habitat (Hayssen 2011). It is tempting to suggest that the reduced requirement to search for food in zoos coupled with the energy-dense diet is responsible for this inactivity. It would be interesting to test whether changing zoo sloth diets towards rations higher in fibre and lower in energy density, as propagated by Bissell (2021), leads to increased activity and hence also to a greater display effect and educative value of sloths.

#### Hedgehog *Erinaceus europaeus*

During a single night, hedgehogs must fill their stomach about twice to meet their energy requirements (Yalden 1976). To do so, they move between one and two kilometres per night and spend approximately four hours foraging (Riber 2006). By contrast, a total feeding duration of 22–27 minutes per night was reported for captive individuals (Campbell 1975) and for some captive individuals locomotion stereotypies have been described (Dimelow 1963). Given the large number of hedgehog care centres across Europe, the paucity of newer information on these aspects is surprising. These differences may be representative of other small insectivorous or omnivorous species in captivity, for example armadillos. Comparing reports on nighttime activity in free-ranging (Ancona and Loughry 2009) and zoo armadillos housed in a nighttime enclosure (Kelly and Rose 2024) suggests significantly more foraging and feeding activity in free-ranging specimens.

#### Giant anteater *Myrmecophaga tridactyla*

Similarly to many other animals, giant anteaters spend less time foraging/feeding in zoos than their natural habitat (Bertassoni and Milléo Costa 2010). When the time to complete a meal consisting of typical 'zoo gruel' was recorded, zoo anteaters had 15-minute feeding bouts. This can be compared to data on the feeding bout length for captive animals offered different species of actual termites in the form of broken-up material from termite mounds harvested the previous day, which was between 1 and 7 minutes, and to free-ranging animals that break up a termite mound to feed

themselves, which averages 21 seconds (Redford 1985). This short duration is explained by the fact that in intact termite mounds, 'soldiers' assemble quickly at the site of breakup, outnumbering the 'workers' and defending the mound with their massive jaws causing the anteater to move towards another mound. Mimicking this aspect of natural feeding behaviour would require many small-portion feedings in zoos. This case is an example that important, interesting and plausible biological facts might shape the behaviour and activity budget of free-ranging animals that one would not intuitively think of; reading the biological literature about the animals kept at a zoo hence is important for developing appropriate ideas.

#### Primates

Also in primates, comparisons between free-ranging and zoo animals indicate longer feeding and foraging times in natural habitats (Kamaluddin et al. 2022; Melfi and Feistner 2002). Apart from notes on tigers (De Rouck et al. 2005) and giant otters *Pteronura brasiliensis* (Friedmann et al. 2023), to the authors' knowledge it is especially in primates that a positive effect of captivity on behavioural repertoire has been noted, insofar as the safety and 'spare time' in captivity might facilitate novel positive behaviours not displayed in the wild (van Schaik et al. 2016). Nevertheless, increasing the time spent feeding/foraging is still an important aim in primate husbandry. Yamanashi and Hayashi (2011) showed that when captive chimpanzees *Pan troglodytes* had to work for their food through cognitive tests, they spent about as much daily time acquiring their food this way as the foraging time of free-ranging chimpanzees. By contrast, if they were provided the same amount of food during weekends (when the cognitive test devices were not operating), their feeding time was significantly lower.

Diet choice can have an effect on interactions within a primate group. Feeding a fruit-free diet, i.e. a diet with fewer items that might be perceived as 'worth fighting for' due to their sugar content, has been associated with reduced intragroup aggression or vocalisation (Britt et al. 2015; Viillard et al. 2023).

#### Large carnivores

The last example case touches on a well-known fact: large carnivores do not have 100% hunting success; in other words, when they are motivated to feed they cannot fulfil that motivation instantly but have to try repeatedly. Examples are 20% hunting success in leopards *Panthera pardus* (Balme et al. 2007) and 29% hunting success in lions *Panthera leo* (Balme et al. 2013). These numbers are specific to the respective ecological context of the observed populations and will vary by season, prey and most likely individual predator experience. Hence, they should not be considered 'natural laws' but mainly an indication that a herbivore starting to graze or browse will in most cases be somewhat successful immediately, but a predator may require several attempts to acquire its prey. To the authors' knowledge this has not been replicated in zoos so far—animals are typically 100% sure that they will get the food with which they are presented—regardless of whether access is more or less difficult. To what extent the option to fail and hence the cognitive incentive to focus and the potential of actually 'feeling successful' affects predator behaviour and welfare has to the authors' knowledge not been investigated. Clauss and Schiffmann (2022) propose that the opportunity of failure represents an important aspect of a 'meaningful' life for animals, as an intuitive concept without proof. The only description of a feeding device for zoo carnivores (for cheetah *Acinonyx jubatus*) that withdraws the food if the animals do not pay sufficient attention or make sufficient effort is that of a moving bait zip-line that pulls up the prey after a short period during which it is within reach of the animals (Williams

et al. 1996). However, this study does not explain the overall feeding management, i.e. whether the consequence of a 'failed attempt' is a fasting day or whether the animals will receive the food a bit later anyhow. Designing hygienically sound, escape-safe, technically feasible methods of presenting carnivores with their food (e.g. humanely killed whole prey) in a way that they can periodically fail to obtain the food might represent an attractive next step in carnivore husbandry. Most likely, the experience of failure need not be as frequent as unsuccessful hunting events in the wild to achieve the effect of ensuring full cognitive focus. One possible effect of a task that requires concentrated focus and is experienced as important by the animals might be relaxation afterwards (Krawczel et al. 2005).

## Conclusion

Dietary management in a modern zoo requires more than formulation of an optimal diet in terms of ingredients, brands and supplements. As wild animals in captivity have been removed from their ecological context and the necessity to acquire food on their own, it is important to imitate natural conditions to facilitate feeding and foraging behaviour that is appropriate for the species and hence as naturalistic as feasible under zoo conditions. Not everything can be mimicked: for ethical reasons, it is not acceptable to feed live prey and hence other ways of simulating searching, hunting and prey processing are needed. For logistical reasons, it is difficult to provide plant material such as hay in a way that resembles the natural growth form with the resistance of the roots (and hence 'plucking' or 'pulling off' cannot be simulated). It is the task of managers of zoo animals to find creative alternative solutions.

Active application of knowledge on natural conditions and behaviours is essential for improving animal welfare by dietary and feeding management. Following the four-step model described, one could argue that steps 1–3 should be relatively straightforward and therefore fundamental to guarantee high standards of animal health and welfare in zoological institutions. However, the harsh reality is often more complicated; employment of qualified nutritionists is rare and dietary management is challenging because of zoo management restrictions.

Steps in the development of zoo animal nutrition are not made in synchrony across species or zoos. Captive herbivorous ungulates, for instance, will be provided with a herbivorous diet in most zoological institutions (step 1). Moving forward to steps 2 and 3, zoos tend to find formulation of an appropriate nutrient profile for ungulates more challenging. First, the different grazer-browser foraging strategies (Clauss and Dierenfeld 2008; Clauss et al. 2008; Hofmann 1989) are often not implemented due to real or perceived difficulty in obtaining browse, resulting in zoo diets containing inappropriate types of forage or inappropriate cultivated or commercially available feeds. Second, captive herbivores are almost never exclusively provided with food covering their basic requirements, i.e. roughage supplemented with an appropriate pellet, but are offered produce or even biscuits for training. Many zoos prefer adding particularly well-liked (and hence most likely 'unnatural') items for enrichment rather than following a strategy of providing the normal diet in an 'enriched' way. Constraints in terms of obtaining adequate food resources—either due to a lack of finances for the required staff time or a lack of dedication of such staff—will result in a relapse back to steps 1–2, for example evident in limited availability of browse during winter in temperate zone zoos or inadequate roughage quality.

Whenever a zoo does succeed in the formulation of an optimal zoo diet on which the animal will survive and nutritional and physiological requirements are covered, the step 4 challenge of facilitating natural feeding and foraging behaviour arises. Food provision is often restricted to the operational hours of the zoo

and feeding management of the animals thus may not account for the majority of their day.

To date, as few as 11 EAZA accredited zoos in Europe employ a qualified nutritionist as a staff member to focus on the nutritional needs of the animals in those collections (Samet and Clauss 2023); based on discussions with these specialists, it is evident that they are often still confronted with issues representative of steps 2 and 3 rather than making progress in step 4. By providing the historical outline of steps in the development of zoo animal nutritional management, the authors hope to inspire zoos to assess where they are with respect to the different animal groups in their care. The case study examples are provided to inspire zoos to venture into timely, plausible solutions for step 4—towards making a major contribution to the welfare of animals in their care, not by adding something to an appropriate diet but by presenting the appropriate diet in a way that makes the life of their animals more meaningful.

## Acknowledgments

The authors thank Sally Binding and Holly Farmer for the invitation to contribute to this special issue.

## References

- Allen M.E., Oftedal O.T., Baer D.J. (1996) The feeding and nutrition of carnivores. In: Kleiman D.G., Allen M.E., Thompson K.V., Lumpkin S. (eds.). *Wild Mammals in Captivity: Principles and Techniques*. Chicago, Illinois: University of Chicago Press, 139–147.
- Ancona K.A., Loughry W.J. (2009) Time budgets of wild nine-banded armadillos. *Southeastern Naturalist* 8(4): 587–598. doi:10.1656/058.008.0402
- Balme G., Hunter L., Slotow R. (2007) Feeding habitat selection by hunting leopards *Panthera pardus* in a woodland savanna: Prey catchability versus abundance. *Animal Behaviour* 74(3): 589–598. doi:10.1016/j.anbehav.2006.12.014
- Balme G.A., Batchelor A., de Woronin Britz N., Seymour G., Grover M., Hes L., Macdonald D.W., Hunter L.T.B. (2013) Reproductive success of female leopards *Panthera pardus*: The importance of top-down processes. *Mammal Review* 43(3): 221–237. doi:10.1111/j.1365-2907.2012.00219.x
- Bashaw M.J. (2011) Consistency of captive giraffe behavior under two different management regimes. *Zoo Biology* 30(4): 371–378.
- Bashaw M.J., Kelling A.S., Bloomsmith M.A., Maple T.L. (2007) Environmental effects on the behavior of zoo-housed lions and tigers, with a case study of the effects of a visual barrier on pacing. *Journal of Applied Animal Welfare Science* 10(2): 95–109.
- Bertassoni A., Milléo Costa L.C. (2010) Behavioral repertoire of giant anteater (*Myrmecophaga tridactyla*, Linnaeus 1758) in nature at Serra da Canastra National Park, MG and in captivity at Curitiba Zoo, PR, Brazil. *Revista de Etologia* 9(2): 21–30.
- Bickel A., Geréz L. (1936) Stoffwechselmechanik und Ernährungsart in ihrer Beziehung zur erbgebundenen Stellung des Menschen unter den Herbivoren, Karnivoren und Omnivoren. *Deutsche Medizinische Wochenschrift* 62(41): 1665–1668. doi:10.1055/s-0028-1141345
- Bissell H. (2021) *Sloth Nutrition Guide*. Orlando, Florida: SeaWorld Parks and Entertainment.
- Bloomsmith M.A., Lambeth S.P. (1995) Effects of predictable versus unpredictable feeding schedules on chimpanzee behavior. *Applied Animal Behaviour Science* 44(1): 65–74. doi:10.1016/0168-1591(95)00570-I
- Bond J.C., Lindburg D.G. (1990) Carcass feeding of captive cheetahs (*Acinonyx jubatus*): The effects of a naturalistic feeding program on oral health and psychological well-being. *Applied Animal Behaviour Science* 26(4): 373–382. doi:10.1016/0168-1591(90)90036-D
- Britt A. (1998) Encouraging natural feeding behavior in captive-bred black and white ruffed lemurs (*Varecia variegata variegata*). *Zoo Biology* 17(5): 379–392. doi:10.1002/(SICI)1098-2361(1998)17:5<3C379::AID-ZOO3%3E3.0.CO;2-X
- Britt S., Cowlard K., Baker K., Plowman A. (2015) Aggression and self-directed behaviour of captive lemurs (*Lemur catta*, *Varecia variegata*, *V. rubra* and *Eulemur coronatus*) is reduced by feeding fruit-free diets. *Journal of Zoo and Aquarium Research* 3(2): 52–58. doi:10.19227/jzar.v3i2.119

- Campbell P.A. (1975) Feeding rhythms of caged hedgehogs (*Erinaceus europaeus* L.). *Proceedings of the New Zealand Ecological Society* 22: 14–18.
- Carbone C., Mace G.M., Roberts S.C., Macdonald D.W. (1999) Energetic constraints on the diet of terrestrial carnivores. *Nature* 402: 286–288. doi:10.1038/46266
- Chesney R.W., Hedberg G. (2010) Metabolic bone disease in lion cubs at the London Zoo in 1889: The original animal model of rickets. *Journal of Biomedical Science* 17: S36. doi:10.1186/1423-0127-17-S1-S36
- Clauss M., Dierenfeld E.S. (2008) The nutrition of browsers. In: Fowler M.E., Miller R.E. (eds.). *Zoo and Wild Animal Medicine: Current Therapy* 6. St Louis, Missouri: Saunders Elsevier, 444–454.
- Clauss M., Kienzle E., Hatt J.M. (2003) Feeding practice in captive wild ruminants: Peculiarities in the nutrition of browsers/concentrate selectors and intermediate feeders. A review. In: Fidgett A., Clauss M., Ganslosser U., Hatt J.M., Nijboer J. (eds.). *Zoo Animal Nutrition*. Fürth, Germany: Filander, 27–52.
- Clauss M., Kaiser T., Hummel J. (2008) The morphophysiological adaptations of browsing and grazing mammals. In: Gordon I.J., Prins H.H.T. (eds.). *The Ecology of Browsing and Grazing*. Heidelberg, Germany: Springer, 47–88.
- Clauss M., Lischke A., Botha H., Hatt J.M. (2016) Carcass consumption by domestic rabbits (*Oryctolagus cuniculus*). *European Journal of Wildlife Research* 62: 143–145. doi:10.1007/s10344-015-0980-y
- Clauss M., Müller D.W.H. (2024) Putting zoo animal cancer into perspective. *Zoo Biology* 43(1): 15–21. doi:10.1002/zoo.21802
- Clauss M., Schiffmann C. (2022) Concepts of animal welfare in natural habitats and in zoos: Meaning and anxiety. In: Miller R.E., Calle P.P., Lamberski N. (eds.). *Fowler's Zoo and Wild Animal Medicine: Current Therapy* 10. St Louis, Missouri: Elsevier, 255–260.
- Crissey S.D., Meehan T.P., Langman C., Pruett-Jones M.A. (1999) Vitamin D metabolites 25(OH)D and 1,25(OH)2D and kidney function indices and the relationship to diet in Goeldi's monkeys (*Callimico goeldii*). *Zoo Biology* 18(6): 565–574. doi:10.1002/(SICI)1098-2361(1999)18:6<3C565::AID-ZOO10%3E3.0.CO;2-C
- De Cuyper A., Clauss M., Carbone C., Codron D., Cools A., Hesta M., Janssens G.P.J. (2019) Predator size and prey size—gut capacity ratios determine kill frequency and carcass production in terrestrial carnivorous mammals. *Oikos* 128(1): 13–22. doi:10.1111/oik.05488
- De Rouck M., Kitchener A.C., Law G., Nelissen M. (2005) A comparative study of the influence of social housing conditions on the behaviour of captive tigers (*Panthera tigris*). *Animal Welfare* 14(3): 229–238.
- del Castillo S.M., Bashaw M.J., Patton M.L., Rieches R.R., Bercovitch F.B. (2005) Fecal steroid analysis of female giraffe (*Giraffa camelopardalis*) reproductive condition and the impact of endocrine status on daily time budgets. *General and Comparative Endocrinology* 141(3): 271–281. doi:10.1016/j.ygcen.2005.01.011
- Depauw S., Hesta M., Whitehouse-Tedd K., Vanhaecke L., Verbrugge A., Janssens G.P.J. (2013) Animal fibre: The forgotten nutrient in strict carnivores? First insights in the cheetah. *Journal of Animal Physiology and Animal Nutrition* 97(1): 146–154.
- Depauw S., Verbist L., Stevens J.M.G., Salas M. (2023) Feeding management of giraffe towards positive welfare. *Journal of Zoo and Aquarium Research* 11(4): 400–413. doi:10.19227/jzar.v11i4.769
- Dimelow E.J. (1963) The behaviour of the hedgehog (*Erinaceus europaeus*) in the routine of life in captivity. *Proceedings of the Zoological Society of London* 141(2): 281–289. doi:10.1111/j.1469-7998.1963.tb01613.x
- du Toit J.T., Yetman C.A. (2005) Effects of body size on the diurnal activity budgets of African browsing ruminants. *Oecologia* 143: 317–325. doi:10.1007/s00442-004-1789-7
- Elia J.B., Erb H.N., Houpt K.A. (2010) Motivation for hay: Effects of a pelleted diet on behavior and physiology of horses. *Physiology & Behavior* 101(5): 623–627. doi:10.1016/j.physbeh.2010.09.010
- Ellis A.D., Redgate S., Zinchenko S., Owen H., Barfoot C., Harris P. (2015) The effect of presenting forage in multi-layered haynets and at multiple sites on night time budgets of stabled horses. *Applied Animal Behaviour Science* 171: 108–116. doi:10.1016/j.applanim.2015.08.012
- Fiennes R.N.T.W. (1974) Problems of rickets in monkeys and apes. *Journal of the Royal Society of Medicine* 67(4): 309–314. doi:10.1177/003591577406700446
- Foose T.J. (1982) *Trophic Strategies of Ruminant Versus Nonruminant Ungulates*. Chicago, Illinois: University of Chicago PhD Thesis.
- Friedmann M., Burkevica A., Lagerström L., Gustafsson J., Johnsson J., Lundgren P., Clauss M. (2023) Activity budget and behaviour of giant otters (*Pteronura brasiliensis*) at Parken Zoo, Eskilstuna, Sweden. *Journal of Zoo and Aquarium Research* 11(4): 356–365. doi:10.19227/jzar.v11i4.770
- Gray J. (2017) *Zoo Ethics: The Challenges of Compassionate Conservation*. Ithaca, New York: Comstock Publishing Associates.
- Guerra Aldrigui L., Nogueira-Filho S.L.G., Souza Altino V., Mendes A., Clauss M., da Cunha Nogueira S.S. (2018) Direct and indirect caecotrophy behaviour in paca (*Cuniculus paca*). *Journal of Animal Physiology and Animal Nutrition* 102(6): 1774–1782. doi:10.1111/jpn.12961
- Gull J.M., Stahl M., Osmann C., Ortmann S., Kreuzer M., Hatt J.M., Clauss M. (2015) Digestive physiology of captive giant anteaters (*Myrmecophaga tridactyla*): Determinants of faecal dry matter content. *Journal of Animal Physiology and Animal Nutrition* 99(3): 565–576. doi:10.1111/jpn.12223
- Hayssen V. (2011) *Choloepus hoffmanni* (Pilosa: Megalonychidae). *Mammalian Species* 43(873): 37–55. doi:10.1644/873.1
- Hediger H. (1966) Diet of animals in captivity. *International Zoo Yearbook* 6(1): 37–58. doi:10.1111/j.1748-1090.1966.tb01688.x
- Hiimae K.M. (2000) Feeding in mammals. In: Schwenk K. (ed.). *Feeding Form, Function, and Evolution in Tetrapod Vertebrates*. San Diego, California: Academic Press, 411–448.
- Hofmann R.R. (1989) Evolutionary steps of ecophysiological adaptation and diversification of ruminants: A comparative view of their digestive system. *Oecologia* 78: 443–457. doi:10.1007/BF00378733
- Hume I.D. (2005) Nutrition of marsupials in captivity. *International Zoo Yearbook* 39(1): 117–132. doi:10.1111/j.1748-1090.2005.tb00011.x
- Jeschke J.M. (2007) When carnivores are “full and lazy”. *Oecologia* 152: 357–364. doi:10.1007/s00442-006-0654-2
- Kamaluddin S.N., Matsuda I., Md-Zain B.M. (2022) Activity budget and postural behaviors in orangutans on Bukit Merah Orang Utan island for assessing captive great ape welfare. *Journal of Applied Animal Welfare Science* 25(3): 244–255. doi:10.1080/10888705.2021.1910032
- Kawata K. (2008) Zoo animal feeding: A natural history viewpoint. *Der Zoologische Garten* 78(1): 17–42. doi:10.1016/j.zoolgart.2008.09.004
- Kelly R., Rose P.E. (2024) Assessing the impact of environmental enrichment on behavior in understudied armadillo species: A case study. *Zoo Biology* 43(1): 100–109. doi:10.1002/zoo.21810
- Kleinlugtenbelt C.L.M., Clauss M., Burkevica A., De Cuyper A. (2023) Fasted and furious? Considerations on the use of fasting days in large carnivore husbandry. *Journal of Zoo and Aquarium Research* 11(3): 318–323. doi:10.19227/jzar.v11i3.713
- Krawczel P.D., Friend T.H., Windom A. (2005) Stereotypic behavior of circus tigers: Effects of performance. *Applied Animal Behaviour Science* 95(3–4): 189–198. doi:10.1016/j.applanim.2005.04.010
- Kuhar C.W., Fuller G.A., Dennis P.M. (2013) A survey of diabetes prevalence in zoo-housed primates. *Zoo Biology* 32(1): 63–69. doi:10.1002/zoo.21038
- Lombard L.S., Witte E.J. (1959) Frequency and types of tumors in mammals and birds of the Philadelphia Zoological Garden. *Cancer Research* 19: 127–141.
- Lukas K.E., Hamor G., Bloomsmith M.A., Horton C.L., Maple T.L. (1999) Removing milk from captive gorilla diets: The impact on regurgitation and reingestion (R/R) and other behaviors. *Zoo Biology* 18(6): 515–528.
- Melfi V.A., Feistner A.T.C. (2002) A comparison of the activity budgets of wild and captive Sulawesi crested black macaques (*Macaca nigra*). *Animal Welfare* 11(2): 213–222. doi:10.1017/S0962728600028165
- Mellor D.J. (2015a) Enhancing animal welfare by creating opportunities for positive affective engagement. *New Zealand Veterinary Journal* 63(1): 3–8. doi:10.1080/00480169.2014.926799
- Mellor D.J. (2015b) Positive animal welfare states and encouraging environment-focused and animal-to-animal interactive behaviours. *New Zealand Veterinary Journal* 63(1): 9–16. doi:10.1080/00480169.2014.926800
- Mellor D.J. (2016) Updating animal welfare thinking: Moving beyond the “Five Freedoms” towards “A Life Worth Living”. *Animals* 6(3): 21. doi:10.3390/ani6030021
- Meredith A.L., Prebble J.L. (2017) Impact of diet on faecal output and caecotroph consumption in rabbits. *Journal of Small Animal Practice* 58(3): 139–145. doi:10.1111/jsap.12620
- Mertens D.R. (1997) Creating a system for meeting the fiber requirements of dairy cows. *Journal of Dairy Science* 80(7): 1463–1481. doi:10.3168/jds.S0022-0302(97)76075-2
- Moitié S., Jarvis R., Bandelow S., Byrne S., Dobbs P., Grant M., Reeves C., White K., Liptovszky M., Baiker K. (2022) Vitamin D status in chimpanzees in human care: A Europe wide study. *Scientific Reports* 12: 17625. doi:10.1038/s41598-022-21211-6
- Morfeld K.A., Brown J.L. (2016) Ovarian acyclicity in zoo African elephants (*Loxodonta africana*) is associated with high body condition scores and elevated serum insulin and leptin. *Reproduction, Fertility and*



- Development* 28(5): 640–647. doi:10.1071/RD14140
- Mykytowycz R., Rowley I. (1958) Continuous observations of the activity of the wild rabbit, *Oryctolagus cuniculus* (L.), during 24 hour periods. *CSIRO Wildlife Research* 3(1): 26–31. doi:10.1071/CWR9580026
- Neuhaus P., Ruckstuhl K.E. (2002) The link between sexual dimorphism, activity budgets, and group cohesion: The case of the plains zebra (*Equus burchelli*). *Canadian Journal of Zoology* 80(8): 1437–1441. doi:10.1139/z02-126
- Olivas I., Rodríguez-Latorre A., Estellés F., Calvet S., Villagrà A. (2013) Assessment of activity patterns of growing rabbits in a flux-controlled chamber. *World Rabbit Science* 21(2): 107–110. doi:10.4995/wrs.2013.1187
- Orban D.A., Siegford J.M., Snider R.J. (2016) Effects of guest feeding programs on captive giraffe behavior. *Zoo Biology* 35(2): 157–166. doi:10.1002/zoo.21275
- Pellew R.A. (1984) The feeding ecology of a selective browser, the giraffe (*Giraffa camelopardalis tippelskirchi*). *Journal of Zoology* 202(1): 57–81. doi:10.1111/j.1469-7998.1984.tb04288.x
- Plowman A. (2013) Diet review and change for monkeys at Paignton Zoo Environmental Park. *Journal of Zoo and Aquarium Research* 1(2): 73–77. doi:10.19227/jzar.v1i2.35
- Prebble J.L., Langford F.M., Shaw D.J., Meredith A.L. (2015a) The effect of four different feeding regimes on rabbit behaviour. *Applied Animal Behaviour Science* 169: 86–92. doi:10.1016/j.applanim.2015.05.003
- Prebble J.L., Shaw D.J., Meredith A.L. (2015b) Bodyweight and body condition score in rabbits on four different feeding regimes. *Journal of Small Animal Practice* 56(3): 207–212. doi:10.1111/jsap.12301
- Ratcliffe H.L. (1966) Diets for zoological gardens: Aids to conservation and disease control. *International Zoo Yearbook* 6(1): 4–23. doi:10.1111/j.1748-1090.1966.tb01686.x
- Rechcigl M. (ed.). (1977a) *CRC Handbook Series in Nutrition and Food. Section G: Diets, Culture Media, Food Supplements*. Volume I: Diets for Mammals. Cleveland, Ohio: CRC Press Inc.
- Rechcigl M. (ed.). (1977b) *CRC Handbook Series in Nutrition and Food. Section G: Diets, Culture Media, Food Supplements*. Volume II: Food Habits of, and Diets for Invertebrates and Vertebrates—Zoo Diets. Cleveland, Ohio: CRC Press Inc.
- Redford K.H. (1985) Feeding and food preference in captive and wild giant anteaters (*Myrmecophaga tridactyla*). *Journal of Zoology* 205(4): 559–572. doi:10.1111/j.1469-7998.1985.tb03544.x
- Rees P.A. (2009) Activity budgets and the relationship between feeding and stereotypic behaviors in Asian elephants (*Elephas maximus*) in a zoo. *Zoo Biology* 28(2): 79–97. doi:10.1002/zoo.20200
- Remis M.J. (2002) Food preferences among captive western gorillas (*Gorilla gorilla gorilla*) and chimpanzees (*Pan troglodytes*). *International Journal of Primatology* 23: 231–249. doi:10.1023/A:1013837426426
- Riber A.B. (2006) Habitat use and behaviour of European hedgehog *Erinaceus europaeus* in a Danish rural area. *Acta Theriologica* 51: 363–371. doi:10.1007/BF03195183
- Sadhir S., Eller A.R., Canington S.L., Sholtz S.B. (2022) Investigating factors of metabolic bone disease in baboons (*Papio* spp.) using museum collections. *American Journal of Biological Anthropology* 177(3): 489–500. doi:10.1002/ajpa.24450
- Samet L., Clauss M. (2023) JZAR Special Edition: Proceedings of the 12th European Zoo Nutrition Conference in Riga, Latvia, 20–22 September 2023. *Journal of Zoo and Aquarium Research* 11(4): 350.
- Schmidt D.A., Kerley M.S., Porter J.H., Dempsey J.L. (2005) Structural and nonstructural carbohydrate, fat, and protein composition of commercially available, whole produce. *Zoo Biology* 24(4): 359–373. doi:10.1002/zoo.20057
- Schüßler D., Gürtler W.D., Greven H. (2015) Activity budgets of Rothschild's giraffes (*Giraffa camelopardalis rothschildi* Lydekker, 1903) in the "Zoom Erlebniswelt Gelsenkirchen". *Der Zoologische Garten* 84(1–2): 61–74. doi:10.1016/j.zoolgart.2015.01.002
- Schwitzer C., Kaumanns W. (2001) Body weights of ruffed lemurs (*Varecia variegata*) in European zoos with reference to the problem of obesity. *Zoo Biology* 20(4): 261–269. doi:10.1002/zoo.1026
- Schwitzer C., Polowinsky S.Y., Solman C. (2009) Fruits as foods – Common misconceptions about frugivory. In: Clauss M., Fidgett A.L., Hatt J.M., Huisman T., Hummel J., Janssen G., Nijboer J., Plowman A. (eds.). *Zoo Animal Nutrition IV*. Fürth, Germany: Filander Verlag, 131–168.
- Takahashi N., Suda S., Shinki T., Horiuchi N., Shiina Y., Tanioka Y., Koizumi H., Suda T. (1985) The mechanism of end-organ resistance to 1 $\alpha$ ,25-dihydroxycholecalciferol in the common marmoset. *Biochemical Journal* 227(2): 555–563. doi:10.1042/bj2270555
- Ullrey D.E. (1989) Nutritional wisdom. *Journal of Zoo and Wildlife Medicine* 20(1): 1–2.
- van Schaik C.P., Burkart J., Damerius L., Forss S.I.F., Koops K., van Noordwijk M.A., Schuppli C. (2016) The reluctant innovator: Orangutans and the phylogeny of creativity. *Philosophical Transactions of the Royal Society B* 371(1690): 20150183. doi:10.1098/rstb.2015.0183
- Veasey J.S., Waran N.K., Young R.J. (1996) On comparing the behaviour of zoo housed animals with wild conspecifics as a welfare indicator, using the giraffe (*Giraffa camelopardalis*) as a model. *Animal Welfare* 5(2): 139–153. doi:10.1017/S0962728600018571
- Viallard F., Lefebvre S., Petry A., Vonfeld I., Quintard B. (2023) Multi-criteria study on a change to a fruit-free diet in Cebidae and Cercopithecidae. *Journal of Zoo and Aquarium Research* 11(4): 376–383. doi:10.19227/jzar.v11i4.738
- Wackernagel H. (1966) Feeding wild animals in zoological gardens. *International Zoo Yearbook* 6(1): 23–37. doi:10.1111/j.1748-1090.1966.tb01687.x
- Walldén W. (2023) Giraffe (*Giraffa camelopardalis*) feeding: How stereotypies and other behaviours changed at Kolmården djurpark in Sweden. *Journal of Zoo and Aquarium Research* 11(4): 393–399. doi:10.19227/jzar.v11i4.789
- Williams B.G., Waran N.K., Carruthers J., Young R.J. (1996) The effect of a moving bait on the behaviour of captive cheetahs (*Acinonyx jubatus*). *Animal Welfare* 5(3): 271–281. doi:10.1017/S0962728600018856
- Yalden D.W. (1976) The food of the hedgehog in England. *Acta Theriologica* 21(30): 401–424.
- Yamaguchi A., Kohno Y., Yamazaki T., Takahashi N., Shinki T., Horiuchi N., Suda T., Koizumi H., Tanioka Y., Yoshiki S. (1986) Bone in the marmoset: A resemblance to vitamin D-dependent rickets, type II. *Calcified Tissue International* 39: 22–27. doi:10.1007/BF02555736
- Yamanashi Y., Hayashi M. (2011) Assessing the effects of cognitive experiments on the welfare of captive chimpanzees (*Pan troglodytes*) by direct comparison of activity budget between wild and captive chimpanzees. *American Journal of Primatology* 73(12): 1231–1238. doi:10.1002/ajp.20995
- Ziegler T.E., Kapoor A., Binkley N.C., Rice K.S., Rogers J., Jolly C.J., Phillips-Conroy J.E. (2018) Comparison of vitamin D metabolites in wild and captive baboons. *American Journal of Primatology* 80(12): e22935. doi:10.1002/ajp.22935